

Oral presentation

Open Access

Evidence from in vivo 31 -phosphorus magnetic resonance spectroscopy phosphodiester levels that exhaled ethane is a biomarker of cerebral n-3 polyunsaturated fatty acid peroxidation in humans

Basant K Puri^{*1}, Serena J Counsell¹, Brian M Ross², Gavin Hamilton³, Marcelo G Bustos⁴ and Ian H Treasaden^{1,4}

Address: ¹Imperial College, London, UK, ²Northern Ontario School of Medicine, Lakehead University, Canada, ³University of California, San Diego, USA and ⁴West London Mental Health NHS Trust, UK

* Corresponding author

from International Society on Brain and Behaviour: 3rd International Congress on Brain and Behaviour
Thessaloniki, Greece. 28 November – 2 December 2007

Published: 17 April 2008

Annals of General Psychiatry 2008, **7**(Suppl 1):S83 doi:10.1186/1744-859X-7-S1-S83

This abstract is available from: <http://www.annals-general-psychiatry.com/content/7/S1/S83>

© 2008 Puri et al.; licensee BioMed Central Ltd.

This study tested the hypothesis that exhaled ethane is a biomarker of cerebral n-3 polyunsaturated fatty acid (PUFA) peroxidation in humans. Ethane is released specifically following peroxidation of n-3 PUFAs, probably via: abstraction of a hydrogen of the unsaturated carbon closest to the methyl end; isomerization to a diene radical; addition of oxygen to form a hydroperoxide; and β -scission to a hydroxyl and an alkoxy radical, the latter forming ethane by hydrogen addition. We reasoned that the cerebral source of ethane would be the docosahexaenoic acid component of membrane phospholipids. Breakdown of the latter also releases phosphorylated polar head groups, giving rise to glycerophosphorylcholine and glycerophosphorylethanolamine which can be measured from the 31 -phosphorus neurospectroscopy phosphodiester peak. Schizophrenia patients were chosen because of evidence of increased free radical-mediated damage and cerebral lipid peroxidation in this disorder. Breath samples from eight patients were analyzed using mass spectrometry. Cerebral 31 -phosphorus spectra were obtained from the same patients from 70 70 70 mm³ voxels using an image-selected in vivo spectroscopy pulse sequence. Ethane and percentage phosphodiester levels were positively correlated ($r_s = 0.714$, $p < 0.05$), thus supporting the hypothesis that the measurement of exhaled ethane levels indexes cerebral n-3 lipid peroxidation.